

The Effects of Cations on the Activity of the Gypsy Moth (Lepidoptera: Lymantriidae) Nuclear Polyhedrosis Virus

MARTIN SHAPIRO

Insect Biocontrol Laboratory, USDA-ARS, Beltsville Agricultural Research Center-W, Beltsville, MD 20705

J. Econ. Entomol. 94(1): 1-6 (2001)

ABSTRACT Fourteen cations were tested at a 1% concentration (wt:wt), as chlorides, for their effects on the biological activity of the gypsy moth, *Lymantria dispar* (L.), nuclear polyhedrosis virus (LdMNPV). Cupric chloride was toxic to gypsy moth larvae. Ferrous and ferric chloride were inhibitory to larval growth and development as well as to virus activity. Strontium chloride was inhibitory to virus activity but had no apparent effects on gypsy moth larvae. Six cations had little or no effect on virus activity (i.e., calcium, lanthanum, magnesium, nickel, potassium, sodium), whereas four cations (i.e., cobalt, manganese, ruthenium, zinc) acted as viral enhancers, as indicated by reductions in LC_{50} s.

KEY WORDS *Lymantria dispar*, cations, viral activity, enhancement

SINCE THE DISCOVERY that fluorescent (=optical) brighteners could enhance the biological activity of insect viruses (Hamm and Shapiro 1992; Shapiro and Robertson 1992; Shapiro et al. 1992; Shapiro and Dougherty 1993; Zou and Young 1994, 1996), much effort has been devoted to the mechanism(s) for this enhancement (Adams et al. 1994, Sheppard and Shapiro 1994, Sheppard et al. 1994, Dougherty et al. 1995, Washburn et al. 1998). Currently, the research has provided much valuable data but is primarily in its descriptive stage. The only brighteners that appear to enhance viral activity belong to the class of stilbenes (Shapiro and Dougherty 1993), but not all stilbenes are active (Argauer and Shapiro 1997).

Stilbenes have been studied for many years as inhibitors of human red blood cell exchange (Cabantchik and Rothstein 1972, Rao et al. 1979, Romero and Ortiz 1988, Venglarik et al. 1994, Romero and Boron 1999). These studies have centered on the stilbenes 4,4'-diisothiocyanostilbene-2,2'-disulfonic acid (DIDS) and 4-acetamido-4'-isothiocyanostilbene-2,2'-disulfonic acid (SITS) and related stilbenes as 4,4'-dinitrostilbene-2,2'-disulfonic acid (DNDS) and demonstrated that both anion and calcium transport were inhibited (Niggli et al. 1982, Romero and Ortiz 1988).

The demonstration that stilbenes are inhibitors of calcium transport, led me to investigate whether organic (Nandi et al. 1990, Sunaga et al. 1990, Cartee et al. 1992) and inorganic (Howell 1982, Sigel et al. 1990, Soler et al. 1992) calcium transport inhibitors could act as viral enhancers. This article reports on the effect of inorganic metals on the activity of the gypsy moth, *Lymantria dispar* (L.), nuclear polyhedrosis virus (LdMNPV).

Materials and Methods

Insects and Virus Inocula. The New Jersey strain of *Lymantria dispar* (L.) (USDA-Aphis, Otis Air National Guard Base, MA) was used and insects were reared on a wheat germ-based diet (Bell et al. 1981). The gypsy moth nuclear polyhedrosis virus inoculum (LdMNPV) was the Hamden isolate LDP-226 (U.S. Forest Service, Hamden, CT).

Bioassays. Polyhedral inclusion bodies (PIBs) were decimally diluted in distilled water to produce concentrations ranging from 10^2 to 10^6 PIBs/ml (=0.02–209.6 PIBs/mm²), and 1 ml of inoculum was pipetted onto the surface in each 180-ml container (=4,770 mm²; Sweetheart Cup, Chicago, IL). Ten late second instars (7 d old; average weight, 35 ± 6.8 mg per larva) were placed in each container and were maintained for 21 d at 29°C, 50% RH, and a photoperiod of 12:12 (L:D) h. Tests involving LdMNPV and LdMNPV/cation (1% wt:wt) combinations were repeated five times with 30 larvae per virus dilution per treatment per replicate, 30 untreated larvae per replicate, and 30 larvae per cation treatment per replicate. In a second series of tests, cations that appeared to act as virus enhancers were tested at 0.001, 0.01, 0.10, and 1.00% (wt:wt). These tests were repeated four times.

Cations. The cations used were calcium chloride (CAS #10035-04-8), cobalt chloride (CAS #7791-13-1), cupric chloride (CAS #10125-13-0), ferric chloride (CAS #10025-77-1), ferrous chloride (CAS #13478-10-9), lanthanum chloride (CAS #10025-84-0), magnesium chloride (CAS #7791-20-0), manganese chloride (CAS #13446-34-9), nickel chloride (CAS #7791-20-0), potassium chloride (CAS #7447-40-7), ruthenium red (=ammoniated ruthenium oxychloride; CAS #11103-72-3), sodium chloride (CAS #7647-14-5), strontium chloride (CAS #10025-70-4),

Table 1. Effects of cations on the activity of *Lymantria dispar* nuclearpolyhedrosis virus (LdMNPV) activity: LC₅₀

Treatment ^a	LC ₅₀ (as PIBs/mm ²) ^b (95% CL)	Slope (±SEM)	Activity ratio ^c
LdMNPV alone	9.79 (8.36–11.45)	1.46 (0.01)	1.00
LdMNPV/Monovalent cations			
Potassium	8.36 (6.65–10.55)	1.44 (0.01)	1.17
Sodium	14.03 (10.94–18.01)	1.55 (0.01)	0.71
LdMNPV/Divalent cations			
Calcium	11.64 (9.41–14.38)	1.71 (0.02)	0.82
Cobalt	0.51 (9.41–14.38)	1.30 (0.02)	19.14
Copper	Toxic to larvae		
Iron	>210		<0.05
Magnesium	12.54 (10.29–15.24)	1.66 (0.01)	0.78
Manganese	5.14 (4.09–6.46)	1.06 (0.01)	1.91
Nickel	7.65 (6.25–17.78)	1.22 (0.02)	1.28
Strontium	17.80 (14.21–22.31)	1.53 (0.01)	0.55
Zinc	0.44 (0.34–0.59)	1.05 (0.01)	22.24
LdMNPV/Trivalent cations			
Iron	>210		<0.05
Lanthanum	12.22 (5.01–25.93)	1.67 (0.05)	0.80
Ruthenium	0.05 (0.01–0.23)	1.33 (0.04)	183.14

^a All cations were tested as chlorides; ruthenium was used as the ammoniated oxychloride (ruthenium red). Only cupric chloride was toxic to gypsy moth larvae.

^b LC₅₀s were expressed as PIBs/mm² of surface area; five replicates; five virus concentrations per treatment per replicate; total = 750 larvae per treatment. Gypsy moth NPV (LdMNPV) was diluted either in distilled water (LdMNPV alone) or in aqueous cation solutions (LdMNPV/cation).

^c Activity ratio is calculated by dividing the LC₅₀ for LdMNPV by LC₅₀s for LdMNPV/cation combinations.

and zinc chloride (CAS #7646-85-7) (Sigma, St. Louis, MO). These chemicals were chosen to represent monovalent, bivalent, and trivalent cations and all were obtained as chlorides. In initial tests, all chemicals were used at 1% (wt:wt). In a second series of tests, cobalt, ruthenium red, and zinc were used at 0.001, 0.01, 0.10, and 1.00% (wt:wt).

Statistical Methods. Concentration-mortality and time-mortality regressions were estimated by probit analysis (LeOra Software 1987) to monitor the biological activities of LdMNPV with and without salts. Failure of 95% CL was used as a criterion for significant differences.

Results

Fourteen cations (as chlorides) were tested as enhancers for LdMNPV (Table 1). One cation (i.e., cupric chloride) was toxic to gypsy moth larvae at a 1% concentration and was not tested further. Both ferrous and ferric chloride had detrimental effects on gypsy moth larvae (i.e., stunted) and gypsy moth NPV activity (i.e., greatly increased LC₅₀s). For both iron compounds, LC₅₀s could only be obtained in one replicate and in one replicate no virus-caused mortality was obtained at a concentration of 210 PIBs/mm² (=10⁶ PIBs per container). None of the other cations appeared to have detrimental effects on gypsy moth larvae at the concentration used. Strontium appeared to be the only other cation that inhibited viral activity (Table 1). Potassium, sodium, calcium, magnesium, nickel, and lanthanum had little effect on virus activity. Manganese (twofold), cobalt (19-fold), zinc (22-fold), and ruthenium (180-fold), however, reduced LC₅₀s (Table 1). The addition of cobalt and ruthenium also decreased the speed of kill (i.e., LT₅₀); whereas, the addition of lanthanum and magnesium increased

the LT₅₀s (Table 2). The remaining cations, with the exceptions of copper and iron, appeared to have little effect on the speed of kill (Table 2).

Following this test, a subsequent study was set up to investigate the relationship between cation concentration and virus enhancement, using cobalt, ruthenium, and zinc (Table 3). In general, enhancement activities of these cations increased as the concentra-

Table 2. Effect of cations on LdMNPV activity: LT₅₀

Treatment ^a	LT ₅₀ (95% CL) in days ^b	Relative speed of kill, % ^c
LdMNPV alone	10.9 (10.3–11.5)	—
LdMNPV/Monovalent cation		
Potassium	11.4 (10.8–12.1)	+4.5
Sodium	11.1 (10.3–11.8)	+1.8
LdMNPV/Divalent cation		
Calcium	11.5 (10.9–12.1)	+5.5
Cobalt	9.6 (9.0–10.0)	–11.9
Copper	Toxic to larvae	
Iron	— ^d	
Magnesium	12.3 (11.6–13.0)	+12.8
Manganese	11.7 (10.9–12.5)	+7.3
Nickel	12.1 (11.3–12.9)	+11.0
Strontium	11.6 (11.3–12.3)	+6.4
Zinc	10.8 (10.2–11.5)	–1.0
LdMNPV/Trivalent cation		
Iron	— ^d	
Lanthanum	12.1 (11.6–12.8)	+11.0
Ruthenium	8.9 (8.5–9.4)	–18.3

^a LdMNPV was diluted in distilled water (LdMNPV alone) or in aqueous cation solutions (LdMNPV/cation).

^b LT₅₀s were determined for a virus concentration of 210 PIBs/mm² of surface area and were averaged for five replicates per treatment.

^c The LT₅₀ for LdMNPV alone was used as the standard and all other LT₅₀s were compared to the standard. Negative values indicate that the treatment resulted in faster kill than the standard. Positive values indicate that the treatment resulted in slower kill than the standard.

^d LT₅₀ values could not be determined at a virus concentration of 210 PIBs/mm².

Table 3. Effect of different concentrations of cobalt chloride, ruthenium red, and zinc chloride on LdMNPV activity: LC₅₀

Treatment ^a	Cation concn	LC ₅₀ (PIBs per mm ²) ^b (95% CL)	Slope (±SEM)	Activity ratio ^c
LdMNPV alone	0	8.79 (6.35–12.49)	1.62 (0.01)	1.00
LdMNPV/Cobalt	0.001	7.32 (6.65–10.44)	1.39 (0.01)	1.20
	0.010	4.42 (3.65–5.41)	2.16 (0.03)	1.99
	0.100	2.70 (2.21–3.33)	1.97 (0.02)	3.25
	1.000	0.32 (0.25–0.40)	1.59 (0.01)	27.41
LdMNPV/Ruthenium	0.001	8.07 (6.52–9.96)	2.17 (0.04)	1.09
	0.010	3.90 (3.10–4.91)	1.85 (0.02)	2.25
	0.100	0.22 (0.10–0.50)	1.92 (0.02)	39.94
	1.000	0.03 (0.02–0.08)	1.33 (0.02)	262.13
LdMNPV/Zinc	0.001	13.65 (10.80–16.90)	1.79 (0.03)	0.64
	0.010	5.09 (4.21–6.23)	2.13 (0.05)	1.73
	0.100	3.23 (2.68–3.88)	2.21 (0.05)	2.72
	1.000	0.37 (0.24–0.50)	1.48 (0.02)	23.97

^a LdMNPV was diluted in distilled water (LdMNPV alone) or in an aqueous cation solution (0.001, 0.010, 0.100, 1.00% wt:wt). No larval mortality was observed among larvae exposed to any of the cations.

^b LC₅₀s are expressed as PIBs per mm²; four replicates; five virus concentrations per treatment per replicate; total = 600 larvae per treatment.

^c Activity ratios are calculated by dividing the LC₅₀ for LdMNPV alone by LC₅₀s for each LdMNPV/cation combination.

tion of the cations increased. In the case of cobalt chloride and zinc chloride, the greatest incremental increase in enhancement occurred as the concentration increased from 0.10 to 1.00% (e.g., eight- to nine-fold). For ruthenium red, the greatest increase in enhancement activity (e.g., 18-fold) occurred as the concentration increased from 0.10 to 1.00%. A further sevenfold increase occurred with an increase in ruthenium red concentration to 1.00%, resulting in a total 260-fold increase in LdMNPV activity (Table 3).

The amount of virus required to produce an LC₁₀, LC₃₀, LC₅₀, LC₇₀, and LC₉₀ combinations of LdMNPV/cobalt, LdMNPV/ruthenium, or LdMNPV/zinc was significantly less ($P < 0.05$) than that required for LdMNPV at every point in the mortality curve (Table 4). Whereas, 8.4 PIBs/mm² were required to produce 50% mortality among LdMNPV-treated larvae, only 0.38, 0.34, and .04 PIBs/mm² were required to achieve the same mortality level among LdMNPV/zinc-, LdMNPV/cobalt-, and LdMNPV/ruthenium-treated larvae, respectively. More than 42 PIBs/mm² were required to achieve 90% kill among LdMNPV-treated larvae, whereas only 0.31–2.73 PIBs/mm² were required in the three LdMNPV-cation treatments. At every mortality level, less NPV was needed in the LdMNPV/ruthenium treatment than in any other treatment. In these treatments (e.g., LdMNPV only, LdMNPV/cobalt, LdMNPV/ruthenium, LdMNPV/zinc) the

amount of NPV to increase mortality from 10 to 50% was similar to that required to increase mortality from 50 to 90%. For example, sixfold more NPV was needed to increase virus kill from 10 to 50%, and a further sixfold increase in NPV was needed to increase virus-caused mortality from 50 to 90% among LdMNPV only and LdMNPV/cobalt treatments. Thus, the addition of these cations to LdMNPV suspensions resulted in significant differences in LC₅₀ and LC₉₀ among treatments but did not change the ratios of LC₉₀ to LC₅₀ within a given treatment.

Discussion

Little is known about the effect of cations on viral activity. More than 40 yr ago, research was conducted in Europe and Japan on the induction of virus diseases (Vago 1951, Yamafuji 1952, Aruga 1958, Steinhaus 1958) but the results were often contradictory. Subsequent research on the effects of chemical stressors on virus incidence led to both negative (Bird 1955, Steinhaus 1958, Shapiro 1961) and positive (Yadava 1971, Shapiro and Bell 1982) results, depending on the insect species and the chemical used. In the 1970s and 1980s some effort was made to explain inactivation of NPVs on cotton leaves. McLeod et al. (1977) noted that cotton dew had a pH above 10 and implicated magnesium ions in cotton dew as responsible for the inactivation of *Helicoverpa* (*Heliothis*) *zea* (Boddie)

Table 4. Comparison of LC values for LdMNPV alone and LdMNPV/cobalt chloride (1%), LdMNPV/ruthenium red (1%), and LdMNPV/zinc chloride (1%) combinations

Treatment	LC values (95% CL) as PIBs/mm ^{2a}				
	LC ₁₀	LC ₃₀	LC ₅₀	LC ₇₀	LC ₉₀
LdMNPV alone	1.43 (1.11–1.74)	4.17 (3.53–4.86)	8.79 (7.59–10.17)	18.53 (15.87–34.40)	54.54 (44.23–69.43)
LdMNPV/Cobalt	0.05 (0.03–0.07)	0.15 (0.12–0.19)	0.32 (0.25–0.40)	0.69 (0.53–0.88)	2.05 (1.45–2.91)
LdMNPV/Ruthenium	0.004 (0.001–0.006)	0.014 (0.01–0.02)	0.03 (0.02–0.08)	0.08 (0.04–0.13)	0.31 (0.14–0.69)
LdMNPV/Zinc	0.05 (0.03–0.08)	0.21 (0.14–0.40)	0.37 (0.24–0.50)	0.83 (0.52–1.05)	2.70 (1.79–3.35)

^a LdMNPV was diluted in distilled water (LdMNPV alone) or in aqueous cation solutions (1%). Table 4 represents an expansion of Table 3 to include several LC values.

nuclear polyhedrosis virus HzSNPV. Young et al. (1977) observed that viral inactivation of cotton was much greater than on soybean leaves and that greater amounts of Mg^{2+} were found in cotton dew than in soybean dew. Elleman and Entwistle (1982, 1985a) noted that the NPV from *Spodoptera littoralis* (Boisduval) (SLMNPV) was also inactivated on cotton leaves. Whereas divalent cations in leaf gland exudates were shown to decrease the solubility of PIBs, Mg^{2+} alone was not responsible for the inactivation of the virus (Elleman and Entwistle 1985b).

In the case of the gypsy moth, cupric chloride (1%) was toxic and was not considered further. Both ferrous and ferric chloride were inhibitory to larval growth and development, as well as to the gypsy moth NPV at the 1% concentration. At concentrations of 0.001, 0.01, and 0.10% little or no adverse effects were observed. Strontium chloride (at 1%) had no adverse effect on larvae but appeared to inhibit NPV, as evidenced by an increase in the LC_{50} (Table 1). The great majority of cations, (i.e., monovalent cations as potassium, sodium; divalent cations as calcium, magnesium, and nickel); and trivalent cations as lanthanum), however, had little or no effect (at 1%) on gypsy moth larvae or NPV-caused mortality.

At a 1% concentration, three divalent (cobalt, manganese, and zinc) and one trivalent (ruthenium) cation decreased LC_{50} s of LdMNPV. Manganese enhanced activity by twofold, cobalt and zinc by 20-fold, and ruthenium by 260-fold (Table 1). In all cases, no enhancement occurred at a concentration of 0.001% but became significant at 0.01% (Table 3). In the case of both cobalt and zinc chloride, the greatest increase (i.e., eight- to ninefold) occurred as the concentration was increased from 0.10 to 1.00%. In the case of ruthenium red, the greatest increase (i.e., 18-fold) occurred at lower concentrations (i.e., when the concentration was increased from 0.01 to 0.10%). A further sevenfold increase occurred as the concentration was increased from 0.10 to 1.00%, resulting in a total 260-fold enhancement. In the case of both cobalt and zinc, LC values were reduced by 20- to 30-fold (in comparison to that of LdMNPV alone) at every value from LC_{10} to LC_{90} . In the case of ruthenium red, the greatest effect of ruthenium red (1%) occurred at the LC_{10} , where the value for MNPV/ruthenium red was 380-fold less than that for LdMNPV alone (Table 4). At subsequent LC values (LC_{30} , LC_{50} , LC_{70} , LC_{90}) the differences were reduced but were still highly significant ($P < 0.01$). The effects of these cations on the speed of virus-caused mortality (LT_{50}) were less than those of virus concentration (LC_{50}) (Table 2). The addition of zinc chloride to LdMNPV did not change to LT_{50} , whereas the LT_{50} s were reduced by 1.3 d (=12%), and by 2.0 d (=18%) by the addition of cobalt chloride and ruthenium red, respectively. Using both the lethal concentration (LC_{50}) and the speed of kill (LT_{50}) as criteria of enhancement, ruthenium red appeared to be the most active cation tested.

Although no tests were conducted to investigate the possible mode(s) of action of these enhancing cations, some clues do exist. Several of the cations tested have

been shown to act as calcium channel blockers in different plant and animal systems. Lanthanum (Richard et al. 1990, Schiffman et al. 1995), magnesium (Meissner et al. 1986, Hattori and Maehashi 1994), and nickel (Dryl and Lopatowska 1990, Yamaguchi and Uematsu 1990) have been shown to inhibit calcium uptake or efflux but they did not act as enhancers for LdMNPV; whereas, cobalt (Scheenen et al. 1994, Philosoph-Hadas 1996 et al.), manganese (Yamaguchi and Uematsu 1990), ruthenium red (Howell 1982, Taipale et al. 1989), and zinc (Kruse and Poppele 1985, Koohmaraie 1992) enhanced the activity of LdMNPV. In addition, cobalt chloride has been used as stain for neurons in insects (Jagota and Habibulla 1992, Brockmann and Brueckner 1995) and ruthenium red has been used as a stain for polysaccharides (Kuno et al. 1986, Mattila 1989).

These results lead us to ask what the relationship between calcium channel inhibition and viral enhancement is. Several cations that do not act as enhancers do act as calcium channel inhibitors (i.e., lanthanum, magnesium, nickel), but these non-effects could be a function of cation concentration. The enhancing cations have certainly been shown to act as calcium channel blockers in diverse biological systems, but does the inhibition or blocking of calcium influx or efflux result in an increase of viral activity in a given insect-virus system? This important question is the subject of a larger research effort to determine the factors affecting pathogenicity of LdMNPV as influenced by stilbene optical brighteners, anion transport inhibitors, cation transport inhibitors, and carbonic anhydrase inhibitors in the gypsy moth-LdMNPV system.

References Cited

- Adams, J. R., C. A. Sheppard, M. Shapiro, and G. J. Tompkins. 1994. Light and electron microscopic investigations on the histopathology of the midgut of gypsy moth larvae infected with LdMNPV plus a fluorescent brightener. *J. Invertebr. Pathol.* 64: 156-159.
- Argauer, R., and M. Shapiro. 1997. Fluorescence and relative activities of stilbene optical brighteners for the gypsy moth (Lepidoptera: Lymantriidae) baculovirus. *J. Econ. Entomol.* 90: 416-420.
- Aruga, H. 1958. Mechanism of the virus resistance in the silkworm, *Bombyx mori*. V. On the induction of the polyhedral virus by nitrogen mustard. *J. Sericult. Sci. Jpn.* 27: 10-13 (in Japanese with English summary).
- Bell, R. A., C. D. Owens, M. Shapiro, and J. G. R. Tardif. 1981. Development of mass rearing technology, pp. 598-633. In C. C. Doane and M. L. McManus [eds.], *The gypsy moth: research toward integrated pest management*. U.S. Dep. Agric. Tech. Bull. 1584.
- Bird, F. T. 1955. Virus diseases of sawflies. *Can. Entomol.* 87: 124-127.
- Brockmann, A., and D. Breuckner. 1995. Projection pattern of poreplate sensory neurons in honey bee worker, *Apis mellifera* L. (Hymenoptera: Apidae). *Inter. J. Insect Morphol. Embryol.* 24: 405-411.
- Cabantchik, Z. I., and A. Rothstein. 1972. The nature of the membrane sites controlling anion permeability of human

- red blood cells as determined by studies with disulfonic stilbene derivatives. *J. Membr. Biol.* 10: 311–330.
- Cartee, G. D., C. Briggs-Tung, and J. O. Holloszy. 1992. Diverse effects of calcium channel blockers on skeletal muscle glucose transport. *Am. J. Physiol.* 263 (Regulatory Integrative Comp. Physiol.) 32: R70–R75.
- Dougherty, E. M., K. Guthrie, and M. Shapiro. 1995. In vitro effects of fluorescent brightener on the efficacy of occlusion body dissolution and polyhedral-derived virions. *Biol. Control* 5: 383–388.
- Dryl, S., and A. Lopatowska. 1990. Inhibition of potassium-induced ciliary reversal in *Fabrea salina* by inorganic and organic calcium channel blockers. *Acta Protozol.* 29: 173–178.
- Elleman, C. J., and P. F. Entwistle. 1982. A study of glands on cotton responsible for the high pH and cation concentration of the leaf surface. *Ann. Appl. Biol.* 100:553–558.
- Elleman, C. J., and P. F. Entwistle. 1985a. Inactivation of a nuclear polyhedrosis virus on cotton by the substances produced by the cotton leaf surface glands. *Ann. Appl. Biol.* 106: 83–92.
- Elleman, C. J., and P. F. Entwistle. 1985b. The effect of magnesium ions on the solubility of polyhedral inclusion bodies and the possible role in the inactivation of the nuclear polyhedrosis virus of *Spodoptera littoralis* by the cotton leaf exudates. *Ann. Appl. Biol.* 106: 93–100.
- Hamm, J. J., and M. Shapiro. 1992. Infectivity of fall armyworm (Lepidoptera: Noctuidae) nuclear polyhedrosis virus enhanced by a fluorescent brightener. *J. Econ. Entomol.* 85: 2149–2152.
- Hattori, T., and S. Maehashi. 1994. Augmentation of calcium influx by stannous chloride at mouse motor nerve terminals. *Res. Commun. Chem. Pathol. Pharmacol.* 84: 253–258.
- Howell, J. N. 1982. The interaction between ruthenium red and isolated sarcoplasmic reticulum. *Membr. Biochem.* 4: 235–245.
- Jagota, A., and M. Habibulla. 1992. Neuronal maps of the frontal ganglion of the cockroach, *Periplaneta americana*, prepared by heavy metal iontophoresis. *J. Morphol.* 213: 287–294.
- Koohmaraie, M. 1992. The role of calcium-dependent proteases (calpains) in post mortem proteolysis and meat tenderness. *Biochimie (Paris)* 74: 239–245.
- Kruse, M. N., and R. E. Poppele. 1985. Electrical “tuning” in muscle spindle receptors. *Brain Res.* 496: 303–306.
- Kuno, T., S. Naito, M. Ohta, N. Kido, H. Ito, and N. Kato. 1986. Staining of O-specific polysaccharide chains of lipopolysaccharides with ruthenium red. *Microbiol. Immunol.* 30: 743–752.
- Mattila, T. 1989. A staining technique to measure capsular polysaccharides of *Staphylococcus aureus* on filter membranes. *J. Microbiol. Methods* 9: 323–332.
- McLeod, P. J., W. Yearian, and S. Y. Young. 1977. Inactivation of *Baculovirus heliothis* by ultraviolet irradiation, dew, and temperature. *J. Invertebr. Pathol.* 30: 237–241.
- Meissner, G., E. Darling, and J. Eveleth. 1986. Kinetics of rapid calcium release by sarcoplasmic reticulum: Effects of calcium, magnesium, and adenine nucleotides. *Biochemistry* 25: 236–244.
- Nandi, J., L. King, D. S. Kaplan, and R. A. Levine. 1990. Mechanisms of gastric proton pump inhibition by calcium channel antagonists. *J. Pharmacol. Exp. Therap.* 252: 1102–1107.
- Niggli, V., E. Sigel, and E. Carafoli. 1982. Inhibition of the purified and reconstituted calcium pump of erythrocytes by um levels of DIDS and NAP-taurine. *FEBS Lett.* 138: 164–166.
- Philosoph-Hadas, S., S. Meir, I. Rosenberger, and A. H. Halvay. 1996. Regulation of the gravitropic response and ethylene biosynthesis in gravistimulated snapdragon spikes by calcium chelators and ethylene inhibitors. *Plant Physiol. (Rockville)* 110: 301–310.
- Rao, A., P. Martin, R.A.F. Reithmeier, and L. C. Cantley. 1979. Location of the disulfonic stilbene binding site of the human erythrocyte anion exchange system by fluorescent energy transfer. *Biochemistry* 18: 4505–4516.
- Richard, D. S., S. W. Applebaum, and L. I. Gilbert. 1990. Allatostatic regulation of juvenile hormone production in vitro by the ring gland of *Drosophila melanogaster*. *Mol. Cell. Endocrinol.* 68: 153–162.
- Romero, M. F., and W. F. Boron. 1999. Electrogenic Na⁺/HCO₃⁻ cotransporters: cloning and physiology. *Annu. Rev. Physiol.* 61: 699–723.
- Romero, P. J., and C. E. Ortiz. 1988. Electrogenic behavior of the human red cell Ca²⁺ pump revealed by disulfonic stilbenes. *J. Membr. Biol.* 101: 237–246.
- Scheenen, W. J., H. P. DeKoning, B. G. Jenks, H. Vaudry, and E. W. Roubos. 1994. The secretion of alpha-MSH from *Xenopus melanotropes* involves calcium influx through omega-conotoxin-sensitive voltage-operated calcium channels. *J. Neuroendocrinol.* 6: 457–464.
- Schiffman, S. S., M. S. Suggs, L. Losee, L. A. Gatlin, W. C. Stagner, and R. M. Bell. 1995. Effect of lipid-derived second messengers on electrophysiological taste responses in the gerbil. *Pharmacol. Biochem. Behav.* 52: 49–58.
- Shapiro. 1961. Environmental factors as stresses in a host-parasite relationship involving the wax moth, *Galleria mellonella* L., and the nuclear polyhedrosis virus, *Borrelnavirus* sp. M.S. thesis, Cornell University, Ithaca, NY.
- Shapiro, M., and R. A. Bell. 1982. Enhanced effectiveness of *Lymantria dispar* (Lepidoptera: Lymantriidae) nucleopolyhedrosis virus formulated with boric acid. *Ann. Entomol. Soc. Am.* 75: 346–349.
- Shapiro, M., and E. M. Dougherty. 1993. The use of fluorescent brighteners as activity enhancers for insect pathogenic viruses, pp. 40–46. In R. D. Lumsden and J. L. Vaughn [eds.], *Pest Management: Biologically Based Pest Management*. Proceedings, XVIII BARC Symposium, Beltsville, MD. American Chemical Society Series, Washington, DC.
- Shapiro, M., and J. L. Robertson. 1992. Enhancement of gypsy moth (Lepidoptera: Lymantriidae) baculovirus activity by optical brighteners. *J. Econ. Entomol.* 85: 1120–1124.
- Shapiro, M., J. J. Hamm, and E. M. Dougherty. 1992. Composition and methods for biocontrol using fluorescent brighteners. U.S. Patent 5:124,149.
- Sheppard, C. A., and M. Shapiro. 1994. Physiological and nutritional effect of a fluorescent brightener on nuclear polyhedrosis virus-infected *Lymantria dispar* (L.) larvae (Lepidoptera: Lymantriidae). *Biol. Control* 4: 404–411.
- Sheppard, C. A., M. Shapiro, and J. L. Vaughn. 1994. Reduction of luminal pH in gypsy moth larvae (*Lymantria dispar* L.) following ingestion of nuclear or cytoplasmic polyhedrosis virus/fluorescent brightener on natural and artificial diets. *Biol. Control* 4: 412–420.
- Sigel, E., R. Baur, J. G. Kunkel, and B. Lanzrein. 1990. Demonstration of a voltage-dependent calcium current in follicles of the cockroach *Nauphoeta cinerea*. *Invertebr. Reprod. Dev.* 18: 159–164.
- Soler, F., F. Fernandez-Belda, and J. C. Gomez-Fernandez. 1992. The Ca²⁺ release channel in junctional sarcoplasmic reticulum: gating and blockage by cations. *Int. J. Biochem.* 24: 903–909.

- Steinhaus, E. A. 1958. Stress as a factor in insect disease, pp. 725-730. Proceedings, X International Congress of Entomology, Montreal, Canada. Ent. Soc. Am., College Park, MD.
- Sunanga, K., and M. Ogihara. 1990. Effects of calcium channel blockers and hydralazine on epinephrine-induced stimulation of glucose output from primary cultured rat hepatocytes. *J. Pharmacobiol. Dyn.* 13:653-658.
- Taipale, H. T., R. A. Kauppinen, and H. Komulainen. 1989. Ruthenium red inhibits the voltage-dependent increase in cytosolic free calcium in cortical synaptosomes from guinea pig. *Biochem. Pharmacol.* 38: 1109-1114.
- Vago, C. 1951. Phenomenes de "Latentia" dans une maladie a ultravirus des insectes. *Rev. Can. Biol.* 10: 299-308.
- Venglarik, C. J., A. K. Singh, and R. J. Bridges. 1994. Comparison of nitro-versus-amino 4,4'-substituents of disulfonic stilbenes as chloride channel blockers. *Mol. Cell. Biochem.* 140: 137-146.
- Washburn, J. O., B. A. Kirkpatrick, E. Haas-Stapleton, and L. E. Volkman. 1998. Evidence that the stilbene-derived optical brightener M₂R enhances *Autographa californica* M nucleopolyhedrovirus infection of *Trichoplusia ni* and *Heliothis virescens* by preventing sloughing of infected midgut epithelial cells. *Biol. Control* 11: 58-69.
- Yadaya, R. L. 1971. On the chemical stressors of nuclear-polyhedrosis virus of gypsymoth, *Lymantria dispar* L. *Z. Angew. Entomol.* 69:303-311.
- Yamafuji, K. 1952. Mechanism of artificial virus formation in silkworm tissue. *Enzymologia* 115: 223-231.
- Yamaguchi, M., and T. Uematsu. 1990. Effects of various metals on hepatic bile calcium excretion in rats: The stimulatory effect of zinc is mediated through acetylcholine action. *Res. Exp. Med.* 190: 145-151.
- Young, S. Y., W. Yearian, and S. Y. Kim. 1977. Effect of dew from cotton and soybean foliage on activity of *Heliothis* nuclear polyhedrosis virus. *J. Invertebr. Pathol.* 29: 105-111.
- Zou, Y., and S. Y. Young. 1994. Enhancement of nuclear polyhedrosis virus activity in larval pests of lepidoptera by a stilbene fluorescent brightener. *J. Entomol. Sci.* 29: 130-133.
- Zou, Y., and S. Y. Young. 1996. Use of a fluorescent brightener to improve *Pseudoplusia includens* (Lepidoptera: Noctuidae) nuclear polyhedrosis virus activity in the laboratory and field. *J. Econ. Entomol.* 89: 92-96.

Received for publication 7 October 1998; accepted 20 June 2000.
